

GenJet™ In Vitro DNA Transfection Reagent for Caco-2 Cells (Ver. II)

----- A General Protocol for Transfecting Caco-2 Cells

- 100 µl
 500 µl
 1000 µl



10075 Tyler Place, Suite 19
 Ijamsville, MD 21754
 FAX. 301-560-4919
 TEL. 301-330-5966
 Toll Free. 1-(866)-918-6812
 Email: info@signagen.com
 Web: www.signagen.com

This product is for laboratory research ONLY and not for diagnostic use

Introduction:

GenJet™ In Vitro DNA Transfection Reagent (Ver. II) is upgraded version of GenJet™ In Vitro DNA Transfection Reagent. With a new chemistry, more DNA condensing groups were released in the new version compared with old version GenJet™, leading to 3~20 times more efficient in DNA delivery. GenJet™ (Ver. II) for Caco-2 cells was pre-optimized and conditioned for transfecting Caco-2 cells.

Important Guidelines for Transfection:

- GenJet™ reagent was formulated for DNA transfection ONLY! The following standard protocol is for transfecting Caco-2 cells.
- For better efficiency, choosing a correct protocol is essential. We strongly encourage to use "General Protocol" first. If the "General Protocol" fails to give satisfactory result (e.g., less than 10%), try the "Advanced Protocol" on back page.
- For high efficiency and lower toxicity, transfect cells at high density. ~90% confluency is highly recommended.
- To lower cytotoxicity, transfect cells in presence of serum (10%) and antibiotics.

PART I. A General Protocol for Transfecting Caco-2 Cells:

Step I. Cell Seeding:

Cells should be plated 18 to 24 hours prior to transfection so that the monolayer cell density reaches to the optimal ~90% confluency at the time of transfection. Complete culture medium with serum and antibiotics is freshly added to each well ~60 minutes before transfection.

Table 1. Recommended Amounts for Different Culture Vessel Formats

Culture Dish	Transfection Volume (ml)	Plasmid DNA (µg)	Diluent Volume (mL)	GenJet™ Reagent (µL)
96-well	0.2	0.2	2 x 0.01	0.6
48-well	0.3	0.5	2 x 0.02	1
24-well	0.5	1.0	2 x 0.05	3
6-well	1.0	2	2 x 0.1	6
35 mm dish	1.0	2	2 x 0.1	6
60 mm dish	3	4	2 x 0.25	12
10 cm dish	5	5 - 6	2 x 0.5	15 - 18
T75 flask	6	9 - 12	2 x 0.75	27 - 36
250 ml flask	12	30 - 50	2 x 1.25	90 - 150

Step II. Preparation of GenJet™-DNA Complex and Transfection Procedures

For Caco-2 cells, the optimal ratio of GenJet™ (µL):DNA (µg) is 3:1. To ensure the optimal size of complex particles, we

recommend using serum-free DMEM with High Glucose to dilute DNA and GenJet™ Reagent.

The following protocol is given for transfection in 24-well plates, refer to **Table 1** for transfection in other culture formats. The optimal transfection conditions For Caco-2 cells are given in the standard protocol described below.

- For each well, add 0.5 ml of complete medium with serum and antibiotics freshly ~60 minutes before transfection.
- For each well, dilute 1 µg of DNA into 50 µl of serum-free DMEM with High Glucose. Vortex gently and spin down briefly to bring drops to bottom of the tube.
- For each well, dilute 3 µl of GenJet™ reagent (Ver. II) into 50 µl of serum-free DMEM with High Glucose. Vortex gently and spin down briefly.

Note: Never use Opti-MEM to dilute DNA and GenJet™ reagent as it may disrupt transfection complex.

- Add the diluted GenJet™ Reagent immediately to the diluted DNA solution all at once. **(Important: do not mix the solutions in the reverse order !)**
- Immediately pipette up and down 3~4 times or vortex briefly to mix followed by incubation for ~15 minutes at room temperature to allow GenJet™-DNA complexes to form.

Note: Never keep the DNA/GenJet™ complex longer than 20 minutes

- Add the 100 µl GenJet™/ DNA complex drop-wise onto the medium in each well and homogenize the mixture by gently swirling the plate.
- Remove DNA/GenJet™ complex-containing medium and replace with fresh complete serum/antibiotics containing medium ~24 hours post transfection.
- Check transfection efficiency 24 to 48 hours post transfection.

Storage: GenJet™ DNA In Vitro Transfection Reagent is stable for up to 12 months at +4 °C. This item shipped at ambient temperature

