

GenMute™ siRNA & DNA Transfection Reagent



10075 Tyler Place, Suite 19
Ijamsville, MD 21754
FAX: 301-560-4919
TEL: 301-330-5966
Toll Free: 1-(866)-918-6812
Email: info@signagen.com
Web: www.signagen.com

----- A General protocol for transfecting mammalian cells

- 100 μ l
- 500 μ l
- 1000 μ l

This product is for laboratory research ONLY and not for diagnostic use

Introduction:

GenMute™ Reagent is a novel biodegradable polymer based siRNA and DNA transfection reagent. With our proprietary pH Dependent Conformational Change (PDCC) technology, the biodegradable polymer was chemically modified by addition of pre-screened hydrophobic groups to side chain, making GenMute™ Reagent a versatile and most powerful gene delivery tool. GenMute™ Reagent have been validated to effectively and reproducibly transfect single siRNA, DNA or co-transfect DNA/siRNA to variety of mammalian cells.

Important Guidelines for Transfection:

- GenMute™ reagent was formulated as a versatile gene delivery tool. While this protocol gives procedures for transfecting DNA to mammalian cells, the protocol for siRNA or siRNA/DNA transfection can be obtained from our website.
- For maximum efficiency, we recommend using GenMute™ Buffer to dilute DNA and GenMute™ Reagent. Alternatively serum free RPMI 1640 medium is acceptable as diluent. Never use serum free DMEM which may contain sodium pyruvate and Opti-MEM which contains serum.

Procedures fro Transfecting DNA to Mammalian Cells

1. Cell Seeding:

Cells should be plated 18 to 24 hours prior to transfection so that the monolayer cell density reaches to the optimal ~80% confluency at the time of transfection. Complete culture medium with serum and antibiotics is freshly added to each well 30~60 minutes before transfection.

Note: GenMute™ is NOT interfered by serum and antibiotics, therefore serum and antibiotic containing medium can be used during the entire experiment.

Table 1. A Guideline for DNA Co-transfection Per Cell Culture Vessel

Culture Dish	Growth Medium (ml)	Diluent Volume (μ L)	Plasmid DNA (μ g)	GenMute™ (μ L)
24-well	0.5	50	0.5	1.5
12-well	0.75	75	0.75	2.25
6-well	1.0	100	1.0	3
60 mm	3.0	300	3.0	9
10 cm/flask 75	8.0	800	8.0	24

2. DNA Transfection protocol:

For DNA transfection experiment, we recommend using

0.5 ~ 1.0 μ g DNA per well in a 6-well plate. As a starting point, we recommend using 1.0 μ g DNA and 3.0 μ l GenMute™ reagent per well of a 6-well plate which usually give satisfactory efficiency without visible toxicity.

The following conditions are given per well of a 6 well plate. For other culture format, please refer to **Table 1**.

- For each well, add 1.0 ml of complete medium with serum and antibiotics freshly 30~60 minutes before transfection.
- Dilute 1.0 μ g DNA into 100 μ l of GenMute™ buffer. Mix by pipeting up and down.
- Add 3 μ l GenMute™ reagent, vortex briefly.
- Incubate for ~10 min at RT to let transfection complex form. Never keep the complex longer than 20 minutes.
- Add the transfection complex to the cells drop wise.
- Gently rock the plate back and forth and return the plate to the incubator.
- Replace transfection medium by cell growth medium 24 hours after transfection and analyze as required.

Storage: GenMute™ siRNA & DNA Transfection Reagent is stable for up to 12 months at 4 °C. This item shipped at ambient temperature