

GenJet™ siRNA Transfection Reagent

----- A General protocol for transfecting mammalian cells

- 100 µl
- 500 µl
- 1000 µl



10075 Tyler Place, Suite 19
Ijamsville, MD 21754
FAX. 301-560-4919
TEL. 301-330-5966
Toll Free. 1-(866)-918-6812
Email: info@signagen.com
Web: www.signagen.com

This product is for laboratory research ONLY and not for diagnostic use

Introduction:

GenJet™ Reagent is liposome based siRNA delivery tool which was formulated with our proprietary pH Dependent Conformational Change (PDCC) technology to give efficient and Reproducible gene knockdown on variety of mammalian cells. GenJet™ Reagent have been validated to effectively and reproducibly transfect single siRNA and co-transfect DNA/siRNA to variety of mammalian cells.

Important Guidelines for Transfection:

- GenJet™ transfection reagent was formulated as a siRNA delivery tool. This protocol gives procedures for co-transfecting siRNA/DNA and for transfecting siRNA to mammalian cells.
- For maximum gene silencing, we recommend using GenJet™ Transfection Buffer to dilute siRNA/DNA and GenJet™ Reagent. Never use pyruvate or serum containing mediums like DMEM or Opti-MEM to dilute siRNA/DNA and GenJet™ reagent.

PART I. Standard siRNA Transfection of Adherent Cells

Step I. Preparation of Working Solution of GenJet™ Transfection Buffer:

GenJet™ Transfection Buffer (5x) is provided as 5x concentrated stock solution. To make working solution, dilute one part of the stock solution with 4 parts of ddH₂O into a sterile bottle. The working solution is stable at 4 °C-RT for 12 months.

Step II. Cell Seeding:

Cells should be plated 18 to 24 hours prior to transfection so that the monolayer cell density reaches to the optimal ~50% confluency at the time of transfection. Complete culture medium with serum and antibiotics is freshly added to each well 30-60 minutes before transfection.

Note: GenJet™ reagent is NOT interfered by serum and antibiotics, therefore serum and antibiotic containing medium can be used during the entire experiment.

Table 1. A Guideline for siRNA transfection per cell culture vessel

| Culture Dish | Growth Medium (ml) | Transfection Buffer (µL) | siRNA (pmoles) Final 5.0 or 20 nM | GenJet™ Reagent (µL) |
|------------------|--------------------|--------------------------|-----------------------------------|----------------------|
| 24-well | 0.5 | 50 | 2.5 / <u>10</u> | 1.2 |
| 12-well | 0.75 | 75 | 3.75 / <u>15</u> | 1.9 |
| 6-well | 1.0 | 100 | 5.0 / <u>20</u> | 2.4 |
| 60 mm | 3.0 | 300 | 15 / <u>60</u> | 7.2 |
| 10 cm / Flask 75 | 8.0 | 800 | 40 / <u>160</u> | 20 |

Step III. siRNA Transfection Protocol:

For optimal siRNA-mediated silencing, we recommend using 1-20 nM siRNA. As a starting point, we recommend using 5.0 nM siRNA which usually gives satisfactory silencing result for most adherent cell lines or primary cells. For hard-to-transfection cells, we recommend using a final siRNA concentration of 20 nM. (bold & underlined in **Table 1**). Due to the exceptional siRNA condensing capacity of GenJet™ reagent, we recommend using same amount of GenJet™ reagent for final 1-20 nM of siRNA (**Table 1**). The following conditions are given per well in a 6 well plate. For other culture format, please refer to **Table 1**.

- For each well, add 1.0 ml of complete medium with serum and antibiotics freshly 30-60 minutes before transfection.
- Dilute 5.0 or **20** pmoles siRNA (final concentration of 5.0 or **20** nM respectively per well) into 100 µl of working solution of GenJet™ Transfection Buffer prepared in **Step I**. Vortex to mix followed by brief spin to bring drops to the bottom of the tube.

Note: For maximum gene silencing, dilute siRNA and GenJet™ reagent with GenJet™ Transfection Buffer (1x).

We strongly suggest reconstituting siRNA stock solution at 5.0 µM, so add 1.0 or 2.0 µl siRNA stock solution per well of 6-well plate to make final 5.0 and 10 nM siRNA respectively.

- Add 2.4 µl GenJet™ reagent, mix by pipetting up and down.
- Incubate for ~15 minutes at RT to let transfection complex form.
- Note: Never keep the complex longer than 30 minutes.**
- Add the transfection mix to the cells drop wise. Gently rock the plate back and forth and return the plate to CO₂ incubator.
- Gene silencing is usually measured 24-48 hours post transfection.

PART II. A Standard Protocol for DNA/siRNA Co-transfection

Step I. Preparation of Working Solution of GenJet™ Transfection Buffer:


GenJet™ Transfection Buffer (5x) is provided as 5x concentrated stock solution. To make working solution, dilute one part of the stock solution with 4 parts of

GenJet™ siRNA & DNA Transfection Reagent



10075 Tyler Place, Suite 19
 Ijamsville, MD 21754
 FAX: 301-560-4919
 TEL: 301-330-5966
 Toll Free: 1-(866)-918-6812
 Email: info@signagen.com
 Web: www.signagen.com

----- A General protocol for transfecting mammalian cells



This product is for laboratory research ONLY and not for diagnostic use

ddH₂O into a sterile bottle. The working solution is stable at 4 °C-RT for 12 months.

Step II. Cell Seeding:

Cells should be plated 18 to 24 hours prior to transfection so that the monolayer cell density reaches to the optimal 60-70% confluency at the time of transfection. Complete culture medium with serum and antibiotics is freshly added to each well 30-60 min before transfection.

Note: GenJet™ reagent is NOT interfered by serum and antibiotics, therefore serum and antibiotic containing medium can be used during the entire experiment.

Table 2. A Guideline for DNA & siRNA Co-transfection Per Cell Culture Vessel

| Culture Dish | Growth Medium (ml) | Transfection Buffer (µL) | Plasmid DNA (µg) | siRNA (pmols) Final 5.0 nM | GenJet™ Reagent (µL) |
|------------------|--------------------|--------------------------|------------------|----------------------------|----------------------|
| 24-well | 0.5 | 50 | 0.25 | 2.5 | 1.5 |
| 12-well | 0.75 | 75 | 0.375 | 3.25 | 2.25 |
| 6-well | 1.0 | 100 | 0.5 | 5.0 | 3 |
| 60 mm | 3.0 | 300 | 1.5 | 15 | 9 |
| 10 cm / flask 75 | 8.0 | 800 | 4.0 | 40 | 24 |

Step III. DNA & siRNA co-transfection protocol:

For DNA/siRNA co-transfection experiment, we recommend using 0.5-0.6 µg DNA and 1-10 nM siRNA per well in a 6-well plate. As a starting point, we recommend using 0.5 µg DNA and 5.0 pmoles siRNA (final concentration 5.0 nM) per well of a 6-well plate which usually give satisfactory knockdown effect.

The following conditions are given per well of a 6 well plate. For other culture format, please refer to [Table 2](#).

- For each well, add 1.0 ml of complete medium with serum and antibiotics freshly 30-60 minutes before transfection.
- Dilute 0.5 µg DNA and 5.0 pmoles siRNA (final 5.0 nM) into 100 µl of working solution of GenJet™ Transfection Buffer. Vortex to mix followed by brief spin to bring drops to the bottom of the tube.

Note: For optimal transfection efficiency and maximum gene silencing, GenJet™ Transfection Buffer is a must for diluting siRNA/DNA and GenJet™ reagent. We strongly suggest preparing siRNA stock solution at 5.0 µM, so add 1.0 µl siRNA stock solution per well of 6-well plate to make final 5.0 nM of siRNA.

- Add 3 µl GenJet™ reagent immediately, mix by pipetting up and down.
- Incubate for ~15 min at RT to let transfection complex form.

Note: Never keep the complex longer than 30 minutes.

- Add the transfection complex to the cells drop wise.
- Gently rock the plate back and forth and return the plate to the incubator.

- Replace transfection medium by cell growth medium ~5 hours after transfection when necessary.
- Gene silencing is usually measured 24-48 hours post transfection.

Storage: GenJet™ siRNA Transfection Reagent is stable for up to 12 months at 4 °C. This item shipped at ambient temperature